

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. Name of the medicinal product

Exemestane 25 mg film-coated tablets  
Equisin

### 2. Qualitative and quantitative composition

Each film-coated tablet contains 25 mg of exemestane.

For a full list of excipients, see section 6.1.

### 3. Pharmaceutical form

Film-coated tablet.

White to off-white, round, compound cup, film-coated tablet with “25” on one side and plain on the other side.

### 4. Clinical particulars

#### 4.1 Therapeutic indications

Exemestane is indicated for the adjuvant treatment of postmenopausal women with oestrogen receptor-positive invasive early breast cancer (EBC), following 2 – 3 years of initial adjuvant tamoxifen therapy.

Exemestane is indicated for the treatment of advanced breast cancer in women with natural or induced postmenopausal status whose disease has progressed following anti-oestrogen therapy. Efficacy has not been demonstrated in patients with oestrogen receptor-negative status.

#### 4.2 Posology and method of administration

##### Posology

##### *Adult and elderly patients*

The recommended dose of exemestane is one 25 mg tablet to be taken once daily preferably after a meal.

In patients with early breast cancer, treatment with exemestane should continue until completion of five years of combined sequential adjuvant hormonal therapy (tamoxifen followed by Exemestane), or earlier if tumour relapse occurs.

In patients with advanced breast cancer, treatment with exemestane should continue until tumour progression is evident.

No dose adjustments are required for patients with hepatic or renal insufficiency (see section 5.2).

### *Paediatric population*

Not recommended for use in children.

### **4.3 Contraindications**

Hypersensitivity to the active substance or other pyrrolidone derivatives or any of the excipients listed in section 6.1.

Exemestane is contraindicated in pre-menopausal women and pregnant or lactating women.

### **4.4 Special warnings and precautions for use**

Exemestane should not be administered to women with pre-menopausal endocrine status. Therefore, whenever clinically appropriate, the post-menopausal status should be ascertained by assessment of LH, FSH and oestradiol levels.

Exemestane should be used with caution in patients with hepatic or renal impairment.

Exemestane tablets contain sucrose and should not be administered to patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency.

Exemestane tablets contain methyl parahydroxybenzoate which may cause allergic reactions (possibly delayed).

Exemestane is a potent oestrogen-lowering agent, and a reduction in bone mineral density (BMD) and an increased fracture rate has been observed following administration (see section 5.1). At the commencement of adjuvant treatment with Exemestane, women with osteoporosis or at risk of osteoporosis should have treatment baseline bone mineral health assessment based on current clinical guidelines and practice. Patients with advanced disease should have their bone mineral density assessed on a case-by-case basis. Although adequate data to show the effects of therapy in the treatment of the bone mineral density loss caused by Exemestane are not available, patients treated with Exemestane should be carefully monitored and treatment for, or prophylaxis of, osteoporosis should be initiated in at-risk patients.

Routine assessment of 25 hydroxy vitamin D levels before the start of aromatase inhibitor treatment should be considered, due to the high prevalence of severe deficiency in women with early breast cancer. Women with Vitamin D deficiency should receive supplementation with Vitamin D.

### **4.5 Interaction with other medicinal products and other forms of interaction**

*In vitro*, evidence showed that the drug is metabolised through cytochrome P450 CYP3A4 and aldoketoreductases (see section 5.2) and does not inhibit any of the major CYP isoenzymes. In a clinical pharmacokinetic study, the specific inhibition of CYP3A4 by ketoconazole showed no significant effects on the pharmacokinetics of exemestane.

In an interaction study with rifampicin, a potent CYP450 inducer, at a dose of 600 mg daily and a single dose of exemestane 25 mg, the AUC of exemestane was reduced by 54% and C<sub>max</sub> by 41%. Since the clinical relevance of this interaction has not been evaluated, the co-administration of

medicinal products, such as rifampicin, anticonvulsants (e.g., phenytoin and carbamazepine) and herbal preparations containing hypericum perforatum (St John's Wort) known to induce CYP3A4 may reduce the efficacy of Exemestane.

Exemestane should be used cautiously with medicinal products that are metabolised via CYP3A4 and have a narrow therapeutic window. There is no clinical experience of the concomitant use of Exemestane with other anticancer medicines.

Exemestane should not be co-administered with oestrogen-containing medicines as these would negate its pharmacological action.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

No clinical data on exposed pregnancies are available with Exemestane. Studies on animals have shown reproductive toxicity (see section 5.3). Exemestane is therefore contraindicated in pregnant women.

##### Breastfeeding

It is unknown whether exemestane is excreted into human milk. Exemestane should not be administered to lactating women.

##### Women of perimenopausal status or child-bearing potential

The physician needs to discuss the necessity of adequate contraception with women who have the potential to become pregnant including women who are perimenopausal or who have recently become postmenopausal, until their postmenopausal status is fully established (see sections 4.3 and 4.4).

#### **4.7 Effects on the ability to drive and use machines**

Exemestane has a moderate influence on the ability to drive and use machines.

Drowsiness, somnolence, asthenia and dizziness have been reported with the use of the exemestane. Patients should be advised that, if these events occur, their physical and/or mental abilities required for operating machinery or driving a car may be impaired.

#### **4.8 Undesirable effects**

Exemestane was generally well tolerated across all clinical studies conducted with Exemestane at a standard dose of 25 mg/day, and undesirable effects were usually mild to moderate.

The withdrawal rate due to adverse events was 7.4% in patients with early breast cancer receiving adjuvant treatment with Exemestane following initial adjuvant tamoxifen therapy. The most commonly reported adverse reactions were hot flashes (22%), arthralgia (18%) and fatigue (16%).

The withdrawal rate due to adverse events was 2.8% in the overall patient population with advanced breast cancer. The most commonly reported adverse reactions were hot flashes (14%) and nausea (12%).

Most adverse reactions can be attributed to the normal pharmacological consequences of oestrogen deprivation (e.g., hot flushes).

The reported adverse reactions from clinical studies and post-marketing experience are listed below by system organ class and frequency.

Frequencies are defined as: Very common ( $\geq 1/10$ ); Common ( $\geq 1/100$  to  $< 1/10$ ); Uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); Rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); Very rare ( $< 1/10,000$ ); Not known (cannot be estimated from the available data).

***Blood and lymphatic system disorders:***

*Very common* Leucopenia(\*\*)

*Common* Thrombocytopenia(\*\*)

*Not known* Lymphocyte count decreased(\*\*)

***Immune system disorders:***

*Uncommon* Hypersensitivity

***Metabolism and nutrition disorders:***

*Common* Anorexia

***Psychiatric disorders:***

*Very common* Depression, insomnia

***Nervous system disorders:***

*Very common* Headache, dizziness

*Common* Carpal tunnel syndrome, paraesthesia

*Rare* Somnolence

***Vascular disorders:***

*Very common* Hot flushes

***Gastrointestinal disorders:***

*Very common* Abdominal pain, nausea

*Common* Vomiting, diarrhoea, constipation, dyspepsia,

***Hepatobiliary disorders:***

*Rare* Hepatitis,<sup>(†)</sup> cholestatic hepatitis<sup>(†)</sup>

***Skin and subcutaneous tissue disorders:***

*Very common* Hyperhidrosis

*Common* Alopecia, rash, urticaria, pruritus

*Rare* Acute generalised exanthematous pustulosis<sup>(†)</sup>

***Musculoskeletal and connective tissue disorders:***

*Very common* Joint and musculoskeletal pain<sup>(\*)</sup>

Common Fracture, osteoporosis

**General disorders and administration site conditions:**

Very common Pain, fatigue

Common Oedema peripheral, asthenia

**Investigations:**

Very common Hepatic enzyme increased, blood bilirubin increased, blood alkaline phosphatase increased

(\*) Includes: arthralgia and less frequently, pain in extremities, osteoarthritis, back pain, arthritis, myalgia and joint stiffness.

(\*\*) In patients with advanced breast cancer thrombocytopenia and leucopenia have been rarely reported. An occasional decrease in lymphocytes has been observed in approximately 20% of patients receiving Exemestane, particularly in patients with pre-existing lymphopenia; however, mean lymphocyte values in these patients did not change significantly over time and no corresponding increase in viral infections was observed. These effects have not been observed in patients treated in early breast cancer studies.

(†) Frequency calculated by rule of 3/X.

The table below presents the frequency of pre-specified adverse events and illnesses in the early breast cancer Study Intergroup Exemestane Study (IES), irrespective of causality, reported in patients receiving trial therapy and up to 30 days after cessation of trial therapy.

<b>Adverse events and illnesses</b>	<b>Exemestane (N = 2249)</b>	<b>Tamoxifen (N = 2279)</b>
Hot flushes	491 (21.8%)	457 (20.1%)
Fatigue	367 (16.3%)	344 (15.1%)
Headache	305 (13.6%)	255 (11.2%)
Insomnia	290 (12.9%)	204 (9.0%)
Sweating increased	270 (12.0%)	242 (10.6%)
Gynaecological	235 (10.5%)	340 (14.9%)
Dizziness	224 (10.0%)	200 (8.8%)
Nausea	200 (8.9%)	208 (9.1%)
Osteoporosis	116 (5.2%)	66 (2.9%)
Vaginal haemorrhage	90 (4.0%)	121 (5.3%)
Other primary cancer	84 (3.6%)	125 (5.3%)
Vomiting	50 (2.2%)	54 (2.4%)
Visual disturbance	45 (2.0%)	53 (2.3%)
Thromboembolism	16 (0.7%)	42 (1.8%)
Osteoporotic fracture	14 (0.6%)	12 (0.5%)
Myocardial infarction	13 (0.6%)	4 (0.2%)

In the IES study, the frequency of ischemic cardiac events in the exemestane and tamoxifen treatment arms was 4.5% versus 4.2%, respectively. No significant difference was noted for any individual cardiovascular event including hypertension (9.9% versus 8.4%), myocardial infarction

(0.6% versus 0.2%) and cardiac failure (1.1% versus 0.7%).

In the IES study, exemestane was associated with a greater incidence of hypercholesterolemia compared with tamoxifen (3.7% versus 2.1%).

In a separate double-blinded, randomised study of postmenopausal women with early breast cancer at low risk treated with exemestane (N=73) or placebo (N=73) for 24 months, exemestane was associated with an average 7-9% mean reduction in plasma HDL-cholesterol, versus a 1% increase on placebo. There was also a 5-6% reduction in apolipoprotein A1 in the exemestane group versus 0-2% for placebo. The effect on the other lipid parameters analysed (total cholesterol, LDL cholesterol, triglycerides, apolipoprotein-B and lipoprotein-a) was very similar in the two treatment groups. The clinical significance of these results is unclear.

In the IES study, gastric ulcer was observed at a higher frequency in the exemestane arm compared to tamoxifen (0.7% versus <0.1%). The majority of patients on exemestane with gastric ulcers received concomitant treatment with non-steroidal anti-inflammatory agents and/or had a prior history.

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Adverse Drug Reaction (ADR)/ Serious Adverse Event (SAE) electronic form linked to the MCAZ database using the following link: <https://primaryreporting.who-umc.org/ZW>.

### **4.9 Overdose**

Clinical trials have been conducted with exemestane given up to 800 mg in a single dose to healthy female volunteers and up to 600 mg daily to postmenopausal women with advanced breast cancer; these dosages were well tolerated. The single dose of exemestane that could result in life-threatening symptoms is not known. In rats and dogs, lethality was observed after single oral doses equivalent respectively to 2000 and 4000 times the recommended human dose on a mg/m<sup>2</sup> basis. There is no specific antidote to overdosage and treatment must be symptomatic. General supportive care, including frequent monitoring of vital signs and close observation of the patient, is indicated.

## **5. Pharmacological properties**

### **5.1 Pharmacodynamic properties**

Pharmacological classification: 9.5.2 Antineoplastic and immunosuppressive medicines- Hormone inhibitors.

#### Mechanism of action

Exemestane is an irreversible, steroidal aromatase inhibitor, structurally related to the natural substrate androstenedione. In post-menopausal women, oestrogens are produced primarily from the conversion of androgens into oestrogens through the aromatase enzyme in peripheral tissues. Oestrogen deprivation through aromatase inhibition is an effective and selective treatment for hormone-dependent breast cancer in postmenopausal women. In postmenopausal women,

exemestane p.o. significantly lowered serum oestrogen concentrations starting from a 5 mg dose, reaching maximal suppression (>90%) with a dose of 10-25 mg. In postmenopausal breast cancer patients treated with the 25 mg daily dose, whole-body aromatization was reduced by 98%.

Exemestane does not possess any progestogenic or oestrogenic activity. A slight androgenic activity, probably due to the 17-hydro derivative, has been observed mainly at high doses. In multiple daily dose trials, exemestane had no detectable effects on adrenal biosynthesis of cortisol or aldosterone, measured before or after the ACTH challenge, thus demonstrating its selectivity with regard to the other enzymes involved in the steroidogenic pathway.

Glucocorticoid or mineralocorticoid replacements are therefore not needed. A non-dose-dependent slight increase in serum LH and FSH levels has been observed even at low doses: this effect is, however, expected for the pharmacological class and is probably the result of feedback at the pituitary level due to the reduction in oestrogen levels that stimulate the pituitary secretion of gonadotropins also in postmenopausal women.

### Clinical efficacy and safety

#### *Adjuvant treatment of early breast cancer*

In a multicentre, randomised, double-blind study (IES), conducted in 4724 postmenopausal patients with oestrogen-receptor-positive or unknown primary breast cancer, patients who had remained disease-free after receiving adjuvant tamoxifen therapy for 2 to 3 years were randomised to receive 3 to 2 years of exemestane (25 mg/day) or tamoxifen (20 or 30 mg/day) to complete a total of 5 years of hormonal therapy.

#### *IES 52-month median follow-up*

After a median duration of therapy of about 30 months and a median follow-up of about 52 months, results showed that sequential treatment with exemestane after 2 to 3 years of adjuvant tamoxifen therapy was associated with a clinically and statistically significant improvement in disease-free survival (DFS) compared with continuation of tamoxifen therapy. Analysis showed that in the observed study period, exemestane reduced the risk of breast cancer recurrence by 24% compared with tamoxifen (hazard ratio 0.76;  $p = 0.00015$ ). The beneficial effect of exemestane over tamoxifen with respect to DFS was apparent regardless of nodal status or prior chemotherapy.

Exemestane also significantly reduced the risk of contralateral breast cancer (hazard ratio 0.57,  $p = 0.04158$ ).

In the whole study population, a trend for improved overall survival was observed for exemestane (222 deaths) compared to tamoxifen (262 deaths) with a hazard ratio of 0.85 (log-rank test:  $p = 0.07362$ ), representing a 15% reduction in the risk of death in favour of exemestane. A statistically significant 23% reduction in the risk of dying (hazard ratio for overall survival 0.77; Wald chi-square test:  $p = 0.0069$ ) was observed for exemestane compared to tamoxifen when adjusting for the pre-specified prognostic factors (i.e., ER status, nodal status, prior chemotherapy, use of HRT and use of bisphosphonates).

## 52-month main efficacy results in all patients (intention to treat population) and oestrogen receptor-positive patients

Endpoint Population	Exemestane Events/N(%)	Tamoxifen Events/N (%)	Hazard Ratio (95% CI)	p-value
<b>Disease-free survival <sup>a</sup></b>				
All patients	354 /2352 (15.1%)	453 /2372 (19.1%)	0.76 (0.67-0.88)	0.00015
ER+ patients	289 /2023 (14.3%)	370 /2021 (18.3%)	0.75 (0.65-0.88)	0.00030
<b>Contralateral breast cancer</b>				
All patients	20 /2352 (0.9%)	35 /2372 (1.5%)	0.57 (0.33-0.99)	0.04158
ER+ patients	18 /2023 (0.9%)	33 /2021 (1.6%)	0.54 (0.30-0.95)	0.03048
<b>Breast cancer-free survival <sup>b</sup></b>				
All patients	289 /2352 (12.3%)	373 /2372 (15.7%)	0.76 (0.65-0.89)	0.00041
ER+ patients	232 /2023 (11.5%)	305 /2021 (15.1%)	0.73 (0.62-0.87)	0.00038
<b>Distant recurrence-free survival <sup>c</sup></b>				
All patients	248 /2352 (10.5%)	297 /2372 (12.5%)	0.83 (0.70-0.98)	0.02621
ER+ patients	194 /2023 (9.6%)	242 /2021 (12.0%)	0.78 (0.65-0.95)	0.01123
<b>Overall survival <sup>d</sup></b>				
All patients	222 /2352 (9.4%)	262 /2372 (11.0%)	0.85 (0.71-1.02)	0.07362
ER+ patients	178 /2023 (8.8%)	211 /2021 (10.4%)	0.84 (0.68-1.02)	0.07569

\* Log-rank test; ER+ patients = oestrogen receptor-positive patients;

Disease-free survival is defined as the first occurrence of local or distant recurrence, contralateral breast cancer, or death from any cause;

<sup>b</sup> Breast cancer-free survival is defined as the first occurrence of local or distant recurrence, contralateral breast cancer or breast cancer death;

<sup>c</sup> Distant recurrence-free survival is defined as the first occurrence of distant recurrence or breast cancer death;

<sup>d</sup> Overall survival is defined as the occurrence of death from any cause.

In the additional analysis for the subset of patients with oestrogen receptor-positive or unknown status, the unadjusted overall survival hazard ratio was 0.83 (log-rank test:  $p = 0.04250$ ), representing a clinically and statistically significant 17% reduction in the risk of dying.

Results from the IES bone substudy demonstrated that women treated with exemestane following 2 to 3 years of tamoxifen treatment experienced moderate reduction in bone mineral density. In the overall study, the treatment-emergent fracture incidence evaluated during the 30-month treatment period was higher in patients treated with exemestane compared with tamoxifen (4.5% and 3.3% correspondingly,  $p = 0.038$ ).

Results from the IES endometrial substudy indicate that after 2 years of treatment, there was a median 33% reduction of endometrial thickness in the exemestane-treated patients compared with no notable variation in the tamoxifen-treated patients. Endometrial thickening, reported at the start of the study treatment, was reversed to normal (<5 mm) for 54% of patients treated with exemestane.

### *IES 87-month median follow-up*

After a median duration of therapy of about 30 months and a median follow-up of about 87 months, results showed that sequential treatment with exemestane after 2 to 3 years of adjuvant tamoxifen therapy was associated with a clinically and statistically significant improvement in DFS compared with continuation of tamoxifen therapy. Results showed that in the observed study period, exemestane significantly reduced the risk of breast cancer recurrence by 16% compared with tamoxifen (hazard ratio 0.84;  $p = 0.002$ ).



Overall, the beneficial effect of exemestane over tamoxifen with respect to DFS was apparent regardless of nodal status or prior chemotherapy or hormonal therapy.

Statistical significance was not maintained in a few sub-groups with small sample sizes. These showed a trend favouring exemestane in patients with more than 9 nodes positive, or previous chemotherapy CMF. In patients with nodal status unknown, previous chemotherapy other, as well as unknown/missing status of previous hormonal therapy a non-statistically significant trend favouring tamoxifen was observed.

In addition, exemestane also significantly prolonged breast cancer-free survival (hazard ratio 0.82,  $p = 0.00263$ ), and distant recurrence-free survival (hazard ratio 0.85,  $p = 0.02425$ ).

Exemestane also reduced the risk of contralateral breast cancer, although the effect was no longer statistically significant in this observed study period (hazard ratio 0.74,  $p = 0.12983$ ). In the whole study population, a trend for improved overall survival was observed for exemestane (373 deaths) compared to tamoxifen (420 deaths) with a hazard ratio of 0.89 (log-rank test:  $p = 0.08972$ ), representing an 11% reduction in the risk of death in favour of exemestane. When adjusting for the pre-specified prognostic factors (i.e., ER status, nodal status, prior chemotherapy, use of HRT and use of bisphosphonates), a statistically significant 18% reduction in the risk of dying (hazard ratio for overall survival 0.82; Wald chi-square test:  $p = 0.0082$ ) was observed for exemestane compared to tamoxifen in the whole study population.

In the additional analysis for the subset of patients with oestrogen receptor-positive or unknown status, the unadjusted overall survival hazard ratio was 0.86 (log-rank test:  $p = 0.04262$ ), representing a clinically and statistically significant 14% reduction in the risk of dying.

Results from a bone sub-study indicate that treatment with exemestane for 2 to 3 years following 3 to 2 years of tamoxifen treatment increased bone loss while on treatment (mean % change from baseline for BMD at 36 months: -3.37 [spine], -2.96 [total hip] for exemestane and -1.29 [spine], -2.02 [total hip], for tamoxifen). However, by the end of the 24-month post-treatment period there were minimal differences in the change in BMD from baseline for both treatment groups, with the tamoxifen arm having slightly greater final reductions in BMD at all sites (mean % change from baseline for BMD at 24 months post-treatment -2.17 [spine], -3.06 [total hip] for exemestane and -3.44 [spine], -4.15 [total hip] for tamoxifen).

All fractures reported during treatment and during follow-up were significantly higher in the exemestane group than on tamoxifen (169 [7.3%] versus 122 [5.2%];  $p = 0.004$ ), but no difference was noted in the number of fractures reported as osteoporotic.

#### *IES 119-month final follow-up*

After a median duration of therapy of about 30 months and a median follow-up of about 119 months, results showed that sequential treatment with exemestane after 2 to 3 years of adjuvant tamoxifen therapy was associated with a clinically and statistically significant improvement in DFS compared with continuation of tamoxifen therapy.

Analysis showed that over the observed study period, exemestane reduced the risk of breast cancer recurrence by 14% compared with tamoxifen (hazard ratio 0.86,  $p = 0.00393$ ). The beneficial effect of exemestane over tamoxifen with respect to DFS was apparent regardless of nodal status or prior chemotherapy.

Exemestane also significantly prolonged breast cancer-free survival (hazard ratio 0.83,  $p < 0.00152$ ), and distant recurrence-free survival (hazard ratio 0.86,  $p = 0.02213$ ). Exemestane also reduced the risk of contralateral breast cancer; however, the effect was no longer statistically significant (hazard ratio 0.75,  $p = 0.10707$ ).

In the whole study population, overall survival was not statistically different between the two groups with 467 deaths (19.9%) occurring in the exemestane group and 510 deaths (21.5%) in the tamoxifen group (hazard ratio 0.91,  $p = 0.15737$ , not adjusted for multiple testing). For the subset of patients with oestrogen receptor-positive or unknown status, the unadjusted overall survival hazard ratio was 0.89 (log-rank test:  $p = 0.07881$ ) in the exemestane group relative to the tamoxifen group.

In the whole study population, a statistically significant 14% reduction in the risk of dying (hazard ratio for OS 0.86; Wald chi-square test:  $p = 0.0257$ ) was observed for exemestane compared with tamoxifen when adjusting for the pre-specified prognostic factors (i.e., ER status, nodal status, prior chemotherapy, use of HRT and use of bisphosphonates).

A lower incidence of other second (non-breast) primary cancers was observed in exemestane-treated patients compared with tamoxifen-only treated patients (9.9% versus 12.4%).

In the main study, which had a median follow-up in all participants of 119 months (0 – 163.94) and median duration of exemestane treatment of 30 months (0 – 40.41), the incidence of bone fractures was reported on 169 (7.3%) patients in the exemestane group compared with 122 (5.2%) patients in the tamoxifen group ( $p = 0.004$ ).

### Efficacy Results From IES in Postmenopausal Women With Early Breast Cancer (ITT)

	No. of Events		Hazard Ratio	p-value
	Exemestane	Tamoxifen	Hazard Ratio	
<b>30-Month Median Treatment and 34.5-Month Median Follow-Up</b>				
Disease-free survival <sup>a</sup>	213	306	0.69 (95% CI: 0.58-0.82)	0.00003
Breast cancer-free survival <sup>b</sup>	171	262	0.65 (95% CI: 0.54-0.79)	<0.00001
Contralateral breast cancer	8	25	0.32 (95% CI: 0.15-0.72)	0.00340
Distant recurrence-free survival <sup>c</sup>	142	204	0.70 (95% CI: 0.56-0.86)	0.00083
Overall survival <sup>d</sup>	116	137	0.86 (95% CI: 0.67-1.10)	0.22962
<b>30-Month Median Treatment and 52-Month Median Follow-Up</b>				
Disease-free survival <sup>a</sup>	354	453	0.77 (95% CI: 0.67-0.88)	0.00015
<b>Efficacy Results From IES in Postmenopausal Women With Early Breast Cancer (ITT)</b>				
	No. of Events		Hazard Ratio	p-value
	Exemestane	Tamoxifen	Hazard Ratio	

Breast cancer-free survival <sup>b</sup>	289	373	0.76 (95% CI: 0.65-0.89)	0.00041
Contralateral breast cancer	20	35	0.57 (95% CI: 0.33-0.99)	0.04158
Distant recurrence-free survival <sup>c</sup>	248	297	0.83 (95% CI: 0.70-0.98)	0.02621
Overall survival <sup>d</sup>	222	262	0.85 (95% CI: 0.71-1.02)	0.07362
<b>30-Month Median Treatment and 87-Month Median Follow-Up</b>				
Disease-free survival <sup>a</sup>	552	641	0.84 (95% CI: 0.75-0.94)	0.002
Breast cancer-free survival <sup>b</sup>	434	513	0.82 (95% CI: 0.72-0.94)	0.00263
Contralateral breast cancer	43	58	0.74 (95% CI: 0.50-1.10)	0.12983
Distant recurrence-free survival <sup>c</sup>	353	409	0.85 ((95% CI: 0.74-0.98)	0.02425
Overall survival <sup>d</sup>	373	420	0.89 (95% CI: 0.77-1.02)	0.08972
<b>30-Month Median Treatment and 119-Month Median Follow-Up</b>				
Disease-free survival <sup>a</sup>	672	761	0.86 (95% CI: 0.77-0.95)	0.00393
Breast cancer-free survival <sup>b</sup>	517	608	0.83 (95% CI: 0.74-0.93)	0.00152
Contralateral breast cancer	57	75	0.75 (95% CI: 0.53-1.06)	0.10707
Distant recurrence-free survival <sup>c</sup>	411	472	0.86 (95% CI: 0.75-0.98)	0.02213
Overall survival <sup>d</sup>	467	510	0.91 (95% CI: 0.81-1.04)	0.15737

CI = confidence interval; IES = Intergroup Exemestane Study; ITT = intention-to-treat.

- Disease-free survival is defined as the first occurrence of local or distant recurrence, contralateral breast cancer or death from any cause.
- Breast cancer-free survival is defined as the first occurrence of local or distant recurrence, contralateral breast cancer or breast cancer death.
- Distant recurrence-free survival is defined as the first occurrence of distant recurrence or breast cancer death.
- Overall survival is defined as the occurrence of death from any cause.

### Treatment of advanced breast cancer

In a randomised peer-reviewed controlled clinical trial, Exemestane at the daily dose of 25 mg has demonstrated a statistically significant prolongation of survival, Time to Progression (TTP), Time to Treatment Failure (TTF) as compared to standard hormonal treatment with megestrol acetate in postmenopausal patients with advanced breast cancer that had progressed following, or during, treatment with tamoxifen either as adjuvant therapy or as first-line treatment for advanced disease.

## 5.2 Pharmacokinetic properties

### Absorption

After oral administration of exemestane tablets, exemestane is absorbed rapidly. The fraction of the dose absorbed from the gastrointestinal tract is high. The absolute bioavailability in humans is unknown, although it is anticipated to be limited by an extensive first-pass effect. A similar effect resulted in an absolute bioavailability in rats and dogs of 5%. After a single dose of 25 mg,

maximum plasma levels of 18 ng/ml are reached after 2 hours. Concomitant intake of food increases the bioavailability by 40%.

#### Distribution

The volume of distribution of exemestane, not corrected for the oral bioavailability, is ca 20000l. The kinetics is linear and the terminal elimination half-life is 24 h. Binding to plasma proteins is 90% and is concentration-independent. Exemestane and its metabolites do not bind to red blood cells.

Exemestane does not accumulate in an unexpected way after repeated dosing.

#### Elimination

Exemestane is metabolised by oxidation of the methylene moiety on the 6 position by CYP3A4 isoenzyme and/or reduction of the 17-keto group by aldoketoreductase followed by conjugation. The clearance of exemestane is ca 500 l/h, not corrected for the oral bioavailability. The metabolites are inactive or the inhibition of aromatase is less than the parent compound. The amount excreted unchanged in urine is 1% of the dose. In urine and faeces, equal amounts (40%) of <sup>14</sup>C-labeled exemestane were eliminated within a week.

#### Special populations

##### *Age*

No significant correlation between the systemic exposure of Exemestane and the age of subjects has been observed.

##### *Renal impairment*

In patients with severe renal impairment ( $CL_{cr} < 30$  ml/min) the systemic exposure to exemestane was 2 times higher compared with healthy volunteers. Given the safety profile of exemestane, no dose adjustment is considered to be necessary.

##### *Hepatic impairment*

In patients with moderate or severe hepatic impairment, the exposure of exemestane is 2-3 fold higher compared with healthy volunteers. Given the safety profile of exemestane, no dose adjustment is considered to be necessary.

### **5.3 Preclinical safety data**

#### Toxicological studies

Findings in the repeat dose toxicology studies in rats and dogs were generally attributable to the pharmacological activity of exemestane, such as its effects on reproductive and accessory organs. Other toxicological effects (on the liver, kidney or central nervous system) were observed only at exposures considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use.

#### Mutagenicity

Exemestane was not genotoxic in bacteria (Ames test), in V79 Chinese hamster cells, in rat

hepatocytes or the mouse micronucleus assay. Although exemestane was clastogenic in lymphocytes *in vitro*, it was not clastogenic in two *in vivo* studies.

#### Reproductive toxicology

Exemestane was embryotoxic in rats and rabbits at systemic exposure levels similar to those obtained in humans at 25 mg/day. There was no evidence of teratogenicity.

#### Carcinogenicity

In a two-year carcinogenicity study in female rats, no treatment-related tumours were observed. In male rats, the study was terminated on week 92, because of early death by chronic nephropathy. In a two-year carcinogenicity study in mice, an increase in the incidence of hepatic neoplasms in both genders was observed at the intermediate and high doses (150 and 450 mg/kg/day). This finding is considered to be related to the induction of hepatic microsomal enzymes, an effect observed in mice but not in clinical studies. An increase in the incidence of renal tubular adenomas was also noted in male mice at the high dose (450 mg/kg/day). This change is considered to be species- and gender-specific and occurred at a dose which represents 63-fold greater exposure than occurs at the human therapeutic dose. None of these observed effects is considered to be clinically relevant to the treatment of patients with exemestane.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

#### *Tablet core*

Polysorbate 80  
Mannitol  
Microcrystalline cellulose  
Povidone  
Crospovidone  
Sodium starch glycolate  
Colloidal anhydrous silica  
Magnesium stearate

#### *Tablet coat*

Hypromellose  
Polyethylene glycol  
Titanium dioxide  
Purified water

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

24 months.

### **6.4 Special precautions for storage**

Store below 30°C.

### **6.5 Nature and contents of the container**

The tablets are packed in a clear PVC/PVDC/Aluminum blister pack.

Pack sizes: 3 x 10 tablets.

### **6.6 Special precautions for disposal and handling**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. APPLICANT**

Equity Pharmaceuticals (Pty) Ltd

100 Sovereign Drive, Route 21 Corporate Park, Nellmapius Drive, Irene

Pretoria, 0157

South Africa

## **8. MANUFACTURERS**

EirGen Pharma Ltd

Westside Business Park, Old Kilmeaden Road

Waterford

Ireland

## **9. REGISTRATION DETAILS**

Zimbabwe registration number: 2024/9.5.2/6554

Zimbabwe category for distribution: Prescription Preparations (P.P.)

## **10. DATE OF REVISION OF TEXT**

March 2024